Pt. 343

- (j) For products containing diphenhydramine citrate identified in $\S 341.12(f)$. Children 2 to under 6 years of age: oral dosage is 9.5 milligrams every 4 to 6 hours, not to exceed 57 milligrams in 24 hours.
- (k) For products containing diphenhydramine hydrochloride identified in § 341.12(g). Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 mg in 24 hours.
- (1) For products containing doxylamine succinate identified in §341.12(h). Children 2 to under 6 years of age: oral dosage is 1.9 to 3.125 milligrams every 4 to 6 hours, not to exceed 18.75 milligrams in 24 hours.
- (m) For products containing phenindamine tartrate identified in §341.12(i). Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 milligrams in 24 hours.
- (n) For products containing pheniramine maleate identified in § 341.12(j). Children 2 to under 6 years of age: oral dosage is 3.125 to 6.25 milligrams every 4 to 6 hours, not to exceed 37.5 milligrams in 24 hours.
- (o) For products containing pyrilamine maleate identified in §341.12(k). Children 2 to under 6 years of age: oral dosage is 6.25 to 12.5 milligrams every 6 to 8 hours, not to exceed 50 milligrams in 24 hours.
- (p) For products containing thonzylamine hydrochloride identified in $\S 341.12(l)$. Children 2 to under 6 years of age: oral dosage is 12.5 to 25 milligrams every 4 to 6 hours, not to exceed 150 milligrams in 24 hours.
- (q) For products containing triprolidine hydrochloride identified in §341.12(m). Children 4 to under 6 years of age: oral dosage is 0.938 milligram every 4 to 6 hours, not to exceed 3.744 milligrams in 24 hours. Children 2 to under 4 years of age: oral dosage is 0.625 milligram every 4 to 6 hours, not to exceed 2.5 milligrams in 24 hours. Infants 4 months to under 2 years of age: oral dosage is 0.313 milligram every 4 to 6 hours, not to exceed 1.252 milligrams in 24 hours.
- (r) For products containing diphenhydramine citrate identified in $\S 341.14(a)(5)$. Children 2 to under 6 years of age: oral dosage is 9.5 milli-

grams every 4 hours, not to exceed 57 milligrams in 24 hours.

(s) For products containing diphenhydramine hydrochloride identified in §341.14(a)(6). Children 2 to under 6 years of age: oral dosage is 6.25 milligrams every 4 hours, not to exceed 37.5 milligrams in 24 hours.

[51 FR 35339, Oct. 2, 1986, as amended at 52 FR 30057, Aug. 12, 1987; 54 FR 8509, Feb. 28, 1989; 57 FR 58376, Dec. 9, 1992; 59 FR 4218, Jan. 28, 1994; 59 FR 29174, June 3, 1994; 59 FR 36051, July 15, 1994]

PART 343—INTERNAL ANALGESIC, ANTIPYRETIC, AND ANTIRHEUMATIC DRUG PROD-UCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

Sec.

343.1 Scope.

343.3 Definitions.

Subpart B—Active Ingredients

343.10 [Reserved]

343.12 Cardiovascular active ingredients.

343.13 Rheumatologic active ingredients.

343.20 [Reserved]

343.22 Permitted combinations of active ingredients for cardiovascular-rheumatologic use.

Subpart C—Labeling

343.50 [Reserved]

343.60 [Reserved]

343.80 Professional labeling.

Subpart D—Testing Procedures

 $343.90\,\,$ Dissolution and drug release testing.

Authority: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 63 FR 56814, Oct. 23, 1998, unless otherwise noted.

Subpart A—General Provisions

§343.1 Scope.

(a) An over-the-counter analgesicantipyretic drug product in a form suitable for oral administration is generally recognized as safe and effective and is not misbranded if it meets each of the conditions in this part in addition to each of the general conditions established in § 330.1 of this chapter.